

A case report of fibroadenoma of the breast with characteristic cystic change on a cut surface

Journal:	<i>SAGE Open Medical Case Reports</i>
Manuscript ID	SOMCR-20-0026.R2
Manuscript Type:	Case Report
Date Submitted by the Author:	15-Apr-2020
Complete List of Authors:	Miyoshi, Kazuya; National Hospital Organization, Fukuyama Medical Center, Breast Surgery Takahashi, Hirotohi; National Hospital Organization, Fukuyama Medical Center, Breast Surgery Inagaki, Masaru; National Hospital Organisation Fukuyama Medical Center, Surgery Watanabe, Jiro; National Hospital Organization, Fukuyama Medical Center, Laboratory of Pathology Yamada, Sohsuke; Kanazawa Medical University, Pathology and Laboratory Medicine; National Hospital Organization, Fukuyama Medical Center, Laboratory of Pathology Mizutani, Kenichi; Kanazawa Ika Daigaku,
Keywords:	Pathology
Additional Keywords:	fibroadenoma, cyst, macroscopic view, cystic fibroadenoma, intraductal fibroadenoma, intraductal fibroadenomatosis
Abstract:	Fibroadenoma with remarkable cystic change is very unusual. Opinions differ as to the interpretation of this lesion. Furthermore, there have been few reports focusing on its macroscopic view. We herein report a case of fibroadenoma in a 43-year-old woman. The patient presented herself to a medical doctor's office due to a rapidly growing breast tumor. Based on a core needle biopsy, a benign lesion was suspected, and the tumor was surgically resected. On a macroscopic study, the cut surface of the tumor revealed a remarkably cystic and well-circumscribed lesion with an intracystic polypoid component. Microscopically, a variety of findings of epithelial and stromal proliferation were observed. This is an interesting case not only because a fibroadenoma with prominent cystic change is unusual but also because the breast tumor showed a characteristically cystic appearance on its macroscopic view.

SCHOLARONE™
Manuscripts

Case report

A case report of fibroadenoma of the breast with characteristic cystic change on a cut surface

**Kazuya Miyoshi¹, Hirotoshi Takahashi¹, Masaru Inagaki², Jiro Watanabe³, Sohsuke Yamada^{3,4},
Kenichi Mizutani^{4*}**

¹Department of Breast Surgery, National Hospital Organization, Fukuyama Medical Center, Hiroshima, Japan; ²Department of Surgery, National Hospital Organization, Fukuyama Medical Center, Hiroshima, Japan; ³Department of Laboratory of Pathology, National Hospital Organization, Fukuyama Medical Center, Hiroshima, Japan; and ⁴Department of Pathology and Laboratory Medicine, Kanazawa Medical University, Ishikawa, Japan.

Short running title: Fibroadenoma with remarkable cystic change

Word count: 780 words (Abstract 137 words); References: 9; and 1 Figure.

*Corresponding author: Kenichi Mizutani, Department of Pathology and Laboratory Medicine, Kanazawa Medical University, 1-1 Daigaku, Uchinada, Kahoku, Ishikawa 920-0293, Japan. Tel: 81-

1
2
3
4
5
6 76-286-2211; and E-mail: kmupalm@gmail.com
7
8
9
10
11

12 **Declaration**

13
14
15 **Ethics approval and consent to participate**

16
17
18 Not applicable.
19
20
21
22
23

24 **Consent for Publication**

25
26
27 Consent was obtained for the publication of this case report.
28
29
30
31
32

33 **Availability of Data and Materials**

34
35
36 The dataset supporting the findings and conclusions of this case report is included within the article.
37
38
39
40
41

42 **Funding**

43
44
45 The authors declare that they received no funding for this report.
46
47
48
49
50

51 **Conflicts of Interests**

52
53
54 The authors declare that they have no conflicts of interests.
55
56
57
58
59
60

Author's contributions

KM, HT, and MI made a substantial contribution to the acquisition of data for the article; and were associated with revision of the paper. JW, SY, and KM made histopathological interpretation and diagnosis; and were associated with drafting or revising the article. KM, HT, MI, JW, SY, and KM approved the version to be published; and agreed to be accountable for the work of the paper.

Acknowledgements

We would like to thank all members who were associated with this case for their expert technical assistance, helpful comment and general support.

Abstract

Fibroadenoma with remarkable cystic change is very unusual. Opinions differ as to the interpretation of this lesion. Furthermore, there have been few reports focusing on its macroscopic view. We herein report a case of fibroadenoma in a 43-year-old woman. The patient presented herself to a medical doctor's office due to a rapidly growing breast tumor. Based on a core needle biopsy, a benign lesion was suspected, and the tumor was surgically resected. On a macroscopic study, the cut surface of the tumor revealed a remarkably cystic and well-circumscribed lesion with an intracystic polypoid component. Microscopically, a variety of findings of epithelial and stromal proliferation were observed. This is an interesting case not only because a fibroadenoma with prominent cystic change is unusual but also because the breast tumor showed a characteristically cystic appearance on its macroscopic view.

Keywords: Fibroadenoma, cystic, macroscopic view, cystic fibroadenoma, intraductal fibroadenoma, intraductal fibroadenomatosis.

Background

Fibroadenoma of the breast is a well-circumscribed benign tumor with epithelial and stromal growth. It usually occurs in young women, with an incidence of 18.5% [1, 2], except for complex fibroadenoma,

1
2
3
4
5
6 which is an unusual variant of fibroadenoma occurring in elderly women [3]. Although cystic changes
7
8
9 are sometimes present, fibroadenoma with predominant cystic change is very unusual [3, 4, 5], and
10
11
12 the interpretation of the lesion seems to be unclear. While the exact incidence of the lesion is unclear,
13
14
15 in 2015, Bhat et al. reported that only one case had been described thus far [3]. When the tumor is
16
17
18 obviously cystic, it resembles an intraductal papilloma [4, 6]. Some papers have stated that
19
20
21 fibroadenoma with remarkable cystic change is called cystic fibroadenoma and classified under
22
23
24 complex fibroadenoma [3, 5, 7]. There are also a few reports describing benign fibroepithelial lesions
25
26
27 with cystic features and intracystic or intraductal components as intraductal fibroadenoma or
28
29
30 intraductal fibroadenomatosis [2, 8, 9]. Most previous reports have focused on the microscopic view
31
32
33 of these lesions.
34

35
36 We herein report an unusual case of fibroadenoma with remarkable cystic change and intracystic
37
38
39 growth showing a characteristic macroscopic view.
40
41
42
43
44

45 **Case presentation**

46
47
48 A 43-year-old woman presented herself to a medical doctor's office due to a rapidly growing tumor
49
50
51 of the right breast. Ultrasonography revealed a middle- to high-echoic mass. A core needle biopsy was
52
53
54 performed showing hypocellular myxoid stroma with small spindle cells and mammary ducts without
55
56
57 atypia (Figure 1A). Based on the core needle biopsy findings, a benign lesion, such as phyllodes tumor
58
59
60

1
2
3
4
5
6 or fibroadenoma, was suspected, and the tumor was surgically resected about two weeks after the
7
8
9 biopsy. On a macroscopic study, the cut surface of the tumor revealed a remarkably cystic and well-
10
11
12 circumscribed lesion with an intracystic polypoid component, measuring 4.0×3.0 cm in diameter with
13
14
15 a whitish to tan color and hemorrhage (Figure 1B). Microscopically, a variety of findings of epithelial
16
17
18 and stromal proliferation, such as a pericanalicular pattern, an intracanalicular pattern, a polypoid or
19
20
21 leaf-like structure with stromal growth, and proliferation of small tubular structures, were observed
22
23
24 (Figure 1C, D). The stromal component had mild to moderate cellularity, showing myxoid change in
25
26
27 some areas. The mitotic activity was low. Infarction and hemorrhage were also observed, but there
28
29
30 were no findings of malignancy.
31
32

33
34 Although phyllodes tumor was considered as a differential diagnosis, she was ultimately
35
36 diagnosed with fibroadenoma (intraductal fibroadenoma) based on these findings.
37
38
39
40
41

42 **Discussion**

43
44
45 The most unique point of this case is the remarkable cystic change in the tumor; however, the cyst was
46
47
48 filled with intracystic polypoid component. Retrospectively, the mass with middle to high echo
49
50
51 intensity on ultrasonography was likely the intracystic component, and the cystic change in the lesion
52
53
54 might be unremarkable.
55
56

57
58 One of the differential diagnoses of an intracystic mass is a phyllodes tumor. It can be
59
60

1
2
3
4
5
6 difficult to distinguish fibroadenoma with an intracanalicular pattern from phyllodes tumor [8]. In the
7
8
9 present case, the microscopic study revealed an intracanalicular pattern or a leaf-like structure with
10
11
12 stromal growth; however, the lesion showed not only interstitial growth but also growth of the
13
14
15 epithelial component, which led to the diagnosis of fibroadenoma. This is because fibroadenoma is
16
17
18 characterized by biphasic proliferation of epithelial and stromal components, on the other hand,
19
20
21 phyllodes tumor mainly shows stromal growth [1]. While carcinoma was another differential diagnosis,
22
23
24 as the tumor had shown rapid growth, there were no malignant findings microscopically. No second
25
26
27 opinion was obtained in this case.
28
29
30
31
32

33 **Conclusion**

34
35
36 This case is interesting not only because fibroadenoma with remarkable cystic change is unusual but
37
38
39 also because the breast tumor showed a characteristically cystic appearance on its macroscopic view.
40
41
42 Opinions differ as to the interpretation of fibroadenoma with prominent cystic change, and another
43
44
45 fibroepithelial lesion may be one of differential diagnoses, so further investigations will be required
46
47
48 to better understand this entity.
49
50
51
52
53

54 **List of Abbreviation Used**

55
56
57 No abbreviation was used
58
59
60

1
2
3
4
5
6
7
8
9
10 **Declaration**

11
12 **Ethics approval and consent to participate**

13
14
15 Not applicable.
16
17
18
19

20
21 **Consent for Publication**

22
23
24 Consent was obtained for the publication of this case report.
25
26
27
28
29

30
31 **Availability of Data and Materials**

32
33 The dataset supporting the findings and conclusions of this case report is included within the article.
34
35
36
37
38

39
40 **Funding**

41
42 The authors declare that they received no funding for this report.
43
44
45
46
47

48
49 **Conflicts of Interests**

50
51 The authors declare that they have no conflicts of interests.
52
53
54
55
56
57
58
59
60

References

1. Thike AA, Brogi E, Harada O, Oyama T, Tse G. Fibroadenoma. In: The WHO classification of tumours editorial board, editors. WHO classification of tumours, 5th edition, breast tumours. Lyon: International agency for research on cancer, 2019, pp. 168-171.
2. Chung A, Scharre K, Wilson M. Intraductal fibroadenomatosis: an unusual variant of fibroadenoma. *The breast journal*. 2008;14(2):193-5.
3. Bhat A, Vijaya C, Gowda VS. Cystic fibroadenoma: report of a rare case with review of literature. *Breast disease*. 2015;35(1):49-52.
4. Rosen PP. Fibroadenoma. *Rosen's breast pathology*, third edition. Lippincott Williams & Wilkins, 2008, pp. 187-202.
5. Durak MG, Karaman I, Canda T, Balci P, Harmancioğlu O. Cystic fibroadenoma of the breast: a case report. *Türk patoloji dergisi*. 2011;27(3):254-6.
6. Cummings MC, da Silva L, Papadimos DJ, Lakhani SR. Fibroadenoma and intraduct papilloma--a common pathogenesis? *Virchows Archiv : an international journal of pathology*. 2009;455(3):271-5.
7. Belmajdoub M, Jayi S. Cystic fibroadenoma detected incidentally in a patient with postpartum infectious mastitis. *The Pan African medical journal*. 2017;28:148.


- 1
2
3
4
5
6
7 8. Hayano F, Yamada S, Nakano S, Watanabe T, Sasaguri Y, Koga S. Intraductal fibroadenoma
8
9 under the nipple in an 11-year-old female. *Diagnostic pathology*. 2014;9:32.
10
11
12 9. Bezić J. Intraductal fibroadenoma and intraductal phyllodes tumour--a part of the spectrum of the
13
14 breast ductal adenoma? *Virchows Archiv : an international journal of pathology*.
15
16 2010;456(1):105-6.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Figure legend

Figure 1. Microscopic findings of the biopsy specimen, and macroscopic and microscopic findings of the surgical specimen. (A) In the biopsy specimen, myxoid stroma with low cellularity and mammary ducts without atypia were observed. (B) The cut surface of the surgical specimen revealed a remarkably cystic and well-circumscribed lesion with an intracystic polypoid component, measuring 4.0×3.0 cm in diameter with a whitish to tan color and hemorrhage. (C, D) On a microscopic study of the surgical specimen, the intracystic component showed a variety of findings of epithelial and stromal proliferation. A pericanalicular pattern, an intracanalicular pattern, a leaf-like structure with stromal growth (C), and proliferation of small tubular structures (D), were observed. (A) Bar = 200 μ m (H&E staining; original magnification: \times 100). (C, D) Bar = 1.0 mm (H&E staining; original magnification: \times 12.5).

Figure 1



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

(1) Ethics approval

Our institution does not require ethical approval for reporting individual cases or case series.

(2) Informed consent

Written informed consent was obtained from the patient(s) for their anonymized information to be published in this article.

For Peer Review