

Journal of Chromatography B Special Issue Contents

Vol. 855, 1 August 2007

31ST ANNUAL MEETING OF THE JAPANESE SOCIETY FOR BIOMEDICAL MASS SPECTROMETRY, NAGOYA, JAPAN, 28-29 SEPTEMBER 2006

PREFACE

New challenge of mass spectrometry technique

by T. Niwa (Nagoya, Japan) and A. Shimizu (Osaka, Japan)

REVIEWS

Tandem time-of-flight (TOF/TOF) mass spectrometry and the curved-field reflectron (Review)

by R.J. Cotter, W. Griffith and C. Jelinek (Baltimore, MD, USA)

A review of quantitative methods for proteomic studies (Review)

by C. Fenselau (College Park, MD, USA)

Chemical proteomics for drug discovery based on compound-immobilized affinity chromatography (Review)

by H. Katayama and Y. Oda (Ibaraki, Japan)

Cardiovascular proteomic analysis (Review)

by T. Suzuki and R. Nagai (Tokyo, Japan)

Application of proteomic technologies to discover and identify biomarkers for excessive alcohol consumption: A review (Review)

by F. Nomura, T. Tomonaga, K. Sogawa, D. Wu and T. Ohashi (Chiba, Japan)

Noninvasive human metabolome analysis for differential diagnosis of inborn errors of metabolism (Review)

by T. Kuhara (Ishikawa, Japan)

Analysis of DNA-bound advanced glycation end-products by LC and mass spectrometry
(Review)

by C. Bidmon, M. Frischmann and M. Pischetsrieder (Erlangen, Germany)

Protein glutathionylation and oxidative stress (Review)

by T. Niwa (Nagoya, Japan)

REGULAR PAPERS

Serum protein profile of rheumatoid arthritis treated with anti-TNF therapy (infliximab)

by T. Takeuchi, T. Nakanishi, Y. Tabushi, A. Hata, T. Shoda, T. Kotani, A. Shimizu, T. Takubo, S. Makino and T. Hanafusa (Osaka, Japan)

Application of a metabolomic method combining one-dimensional and two-dimensional gas chromatography-time-of-flight/mass spectrometry to metabolic phenotyping of natural variants in rice

by M. Kusano, A. Fukushima, M. Kobayashi, N. Hayashi (Kanagawa, Japan), P. Jonsson, T. Moritz (Umea, Sweden), K. Ebana (Ibaraki, Japan) and K. Saito (Kanagawa and Chiba, Japan)

ESI-MS/MS study of acylcarnitine profiles in urine from patients with organic acidemias and fatty acid oxidation disorders

by H. Kobayashi, Y. Hasegawa, M. Endo, J. Purevsuren and S. Yamaguchi (Shimane, Japan)

Analysis of bile acid glutathione thioesters by liquid chromatography/electrospray ionization-tandem mass spectrometry

by K. Mitamura, M. Sogabe, H. Sakanashi, S. Watanabe, T. Sakai, Y. Yamaguchi, T. Wakamiya and S. Ikegawa (Higashi-osaka, Japan)

MALDI-based imaging mass spectrometry revealed abnormal distribution of phospholipids in colon cancer liver metastasis

by S. Shimma (Aichi, Japan), Y. Sugiura (Aichi, Tokyo and Kanagawa, Japan), T. Hayasaka (Aichi, Japan), Y. Hoshikawa, T. Noda (Tokyo, Japan) and M. Setou (Aichi, Tokyo and Kanagawa, Japan)

Simple and quantitative analysis of urinary sulfated tauro- and glycodihydroxycholic acids in infant with cholestasis by electrospray ionization mass spectrometry

by T. Shinka, Y. Inoue, M. Ohse and T. Kuhara (Ishikawa, Japan)

Changes in urinary level and configuration ratio of D-lactic acid in patients with short bowel syndrome

by Y. Inoue, T. Shinka, M. Ohse, M. Kohno, K. Konuma, H. Ikawa and T. Kuhara (Ishikawa, Japan)

Simultaneous determination of 13 amphetamine related drugs in human whole blood using an enhanced polymer column and gas chromatography-mass spectrometry

by K. Kudo, T. Ishida, K. Hara, S. Kashimura, A. Tsuji and N. Ikeda (Fukuoka, Japan)